

Complete Genome Sequence of CHYJ130330, a Highly Virulent Strain of Porcine Epidemic Diarrhea Virus in South China

Aiqing Jia, Xiaosheng Feng, Qi Liu, Ruyue Zhou, Guiping Wang

Guangdong Haid Institute of Animal Husbandry & Veterinary, PanYu, Guangzhou, Guangdong, China

A.J. and X.F. contributed equally to this work.

Porcine epidemic diarrhea virus (PEDV) strain CHYJ130330 was isolated from southern China and shown to be highly virulent when inoculated into neonatal pigs. This report describes the complete genome sequence of CHYJ130330. These data will provide important insights into the variation of PEDV in China.

Received 10 February 2014 Accepted 26 March 2014 Published 17 April 2014

Citation Jia A, Feng X, Liu Q, Zhou R, Wang G. 2014. Complete genome sequence of CHYJ130330, a highly virulent strain of porcine epidemic diarrhea virus in south China. *Genome Announc.* 2(2):e00165-14. doi:10.1128/genomeA.00165-14.

Copyright © 2014 Jia et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Aiqing Jia, ainqing-jia@hotmail.com.

Porcine epidemic diarrhea virus (PEDV) can cause acute diarrhea and dehydration in 2- to 7-day-old piglets. Infected piglets die within 2 to 3 days. The mortality rate associated with this virus is 90% (1). PEDV emerged in the 1980s and has since become more serious, having resulted in severe economic losses in China (2–4).

In March 2013, the wild-type PEDV strain CHYJ130330 was obtained from a fecal specimen of a 3-day-old dead piglet on a commercial swine farm in Guangdong Province, southern China. This strain was successfully passaged in a Vero cell line. Twenty-four hours after inoculation, the Vero cells demonstrated an obvious cytopathic effect characterized by cell fusion and syncytium formation. The CHYJ130330 strain demonstrates high virulence; the oral ingestion of 10 \times the 50% tissue culture infective dose (TCID₅₀) of this virus by 2- to 7-day-old piglets can induce severe diarrhea after 48 h.

To determine the complete-genome sequence of the PEDV isolate CHYJ130330, 21 pairs of primers based on this isolate were designed to generate overlapping amplicons by reverse transcription-PCR. The complete genome of CHYJ130330 is 28,038 nucleotides (nt) in length, with a poly(A) tail. The genomic organization of the isolate is similar to that previously described and includes a 5' untranslated region (UTR) (nt 1 to 292), a replicase gene comprising open reading frame 1a (ORF1a) (nt 293 to 12601) and ORF1b (nt 12601 to 20637), a spike gene (nt 20634 to 24794), ORF3 (nt 24794 to 25468), an envelope gene (nt 25449 to 25679), a membrane gene (nt 25687 to 26367), a nucleoprotein gene (nt 26379 to 27704), and a 3' UTR (nt 27705 to 28038). The complete genome of PEDV strain CHYJ130330 has nucleotide identities of 96.5% to 99.3% with other complete PEDV genomes available in GenBank (1, 2, 5–9). We observed a high nucleotide identity of 99.1% with the U.S. strain IA1 (10), which was identified on 29 April 2013 by the Iowa State University Veterinary Diagnostic Laboratory and corroborated on 16 May 2013 by the National Veterinary Services Laboratories (5). Porcine epidemic diarrhea was recognized in U.S. swine for the first time in early

2013; since then, PEDV has been detected in 18 U.S. states and has resulted in large economic losses (6, 7).

In conclusion, the PEDV isolate CHYJ130330 is a highly virulent strain that may be a candidate for vaccine development and diagnostic techniques. Investigations of its immunogenicity and pathogenicity are ongoing.

Nucleotide sequence accession number. The complete genome sequence of PEDV strain CHYJ130330 has been deposited in GenBank under the accession no. [KJ020932](https://www.ncbi.nlm.nih.gov/nuccore/KJ020932).

ACKNOWLEDGMENT

This work was supported by a grant from the Science and Technology Planning Project of Guangdong Province, China (2012A020602032).

REFERENCES

- Zhao M, Sun Z, Zhang Y, Wang G, Wang H, Yang F, Tian F, Jiang S. 2012. Complete genome sequence of a Vero cell-adapted isolate of porcine epidemic diarrhea virus in eastern China. *J. Virol.* 86:13858–13859. <http://dx.doi.org/10.1128/JVI.02674-12>.
- Chen J, Liu X, Shi D, Shi H, Zhang X, Feng L. 2012. Complete genome sequence of a porcine epidemic diarrhea virus variant. *J. Virol.* 86:3408. <http://dx.doi.org/10.1128/JVI.07150-11>.
- Li ZL, Zhu L, Ma JY, Zhou QF, Song YH, Sun BL, Chen RA, Xie QM, Bee YZ. 2012. Molecular characterization and phylogenetic analysis of porcine epidemic diarrhea virus (PEDV) field strains in south China. *Virus Genes* 45:181–185. <http://dx.doi.org/10.1007/s11262-012-0735-8>.
- Stevenson GW, Hoang H, Schwartz KJ, Burrough ER, Sun D, Madson D, Cooper VL, Pillatzki A, Gauger P, Schmitt BJ, Koster LG, Killian ML, Yoon KJ. 2013. Emergence of porcine epidemic diarrhea virus in the United States: clinical signs, lesions, and viral genomic sequences. *J. Vet. Diagn. Invest.* 25: 649–654. <http://dx.doi.org/10.1177/1040638713501675>.
- Marthaler D, Jiang Y, Otterson T, Goyal S, Rossow K, Collins J. 2013. Complete genome sequence of porcine epidemic diarrhea virus strain USA/Colorado/2013 from the United States. *Genome Announc.* 1(4): e00555-13. <http://dx.doi.org/10.1128/genomeA.00555-13>.
- Fan H, Zhang J, Ye Y, Tong T, Xie K, Liao M. 2012. Complete genome sequence of a novel porcine epidemic diarrhea virus in South China. *J. Virol.* 86:10248–10249. <http://dx.doi.org/10.1128/JVI.01589-12>.
- Hoang H, Killian ML, Madson DM, Arruda PH, Sun D, Schwartz KJ, Yoon KJ. 2013. Full-length genome sequence of a plaque-cloned virulent porcine epidemic diarrhea virus isolate (USA/Iowa/18984/2013)

- from a midwestern U.S. swine herd. *Genome Announc.* 1(6):e01049-13. <http://dx.doi.org/10.1128/genomeA.01049-13>.
8. Chen J, Wang C, Shi H, Qiu HJ, Liu S, Shi D, Zhang X, Feng L. 2011. Complete genome sequence of a Chinese virulent porcine epidemic diarrhea virus strain. *J. Virol.* 85:11538–11539. <http://dx.doi.org/10.1128/JVI.06024-11>.
9. Song D, Park B. 2012. Porcine epidemic diarrhoea virus: a comprehensive review of molecular epidemiology, diagnosis, and vaccines. *Virus Genes* 44:167–175. <http://dx.doi.org/10.1007/s11262-012-0713-1>.
10. Huang YW, Dickerman AW, Pablo P, Li L, Li F, Ross K, Tanja O, Meng XJ. 2013. Origin, evolution, and genotyping of emergent porcine epidemic diarrhea virus strains in the United States. *mBio* 4(5):e00737-13. <http://dx.doi.org/10.1128/mBio.00737-13>.